

On pointwise inner automorphisms of nilpotent groups of class 2

Z. Azhdari, M. Akhavan-Malayeri*;

Department of Mathematics, Alzahra University

Abstract

An automorphism θ of a group G is pointwise inner if $\theta(x)$ is conjugate to x for any $x \in G$. The set of all pointwise inner automorphisms of group G , denoted by $\text{Aut}_{\text{pwi}}(G)$ form a subgroups of $\text{Aut}(G)$ containing $\text{Inn}(G)$. In this paper, we find a necessary and sufficient condition in certain finitely generated nilpotent groups of class 2 for which $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$. We also prove that in a nilpotent group of class 2 with cyclic commutator subgroup $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$ and the quotient $\text{Aut}_{\text{pwi}}(G)/\text{Inn}(G)$ is torsion. In particular if G' is a finite cyclic group then $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$.

MSC: Primary 20D45; Secondary 20E36

Introduction

By definition, a pointwise inner automorphism of a group G is an automorphism $\theta: G \rightarrow G$ such that t and $\theta(t)$ are conjugate for any $t \in G$. This notion appears in the famous book of Burnside [1, Note B, p 463]. Denote by $\text{Aut}_{\text{pwi}}(G)$ the set of all pointwise inner automorphisms of G .

Obviously, $\text{Aut}_{\text{pwi}}(G)$ contains $\text{Inn}(G)$, the group of all inner automorphisms of G . These groups can coincide, for instance when G is $S_n, A_n, \text{SL}_n(D)$ and $\text{GL}_n(D)$ where D is an Euclidean domain (see [7], [10], [11]).

By a result of Grossman [5], it turns out that $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$ when G is a free group. Endimioni in [4] proved that this property remains true in a free nilpotent group.

Keywords: Inner Automorphism, Pointwise Inner Automorphism, Nilpotent Group.

Received: 26 Jan 2013

Revised 30 June 2013

*Corresponding author: mmalayer@alzahra.ac.ir.

Also Yadav in [12] gave a sufficient condition for a finite p -group G of nilpotent class 2 to be such that $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$. But the equality does not hold in general.

In fact, in 1911, Burnside posed the following question: Does there exist any finite group G such that G has a non-inner and pointwise inner automorphism? In 1913, Burnside himself gave an affirmative answer to this question [3]. Indeed, there are many examples of groups admitting a pointwise inner automorphism which is not inner (see, for instance [3], [4], [8], [9], [12] where these groups are besides nilpotent).

Segal also gave a subtle example. He constructed a finitely generated torsion-free nilpotent group G , in which $\text{Aut}_{\text{pwi}}(G)/\text{Inn}(G)$ contains an element of infinite order (see [9]).

In this paper we study the pointwise inner automorphisms of a finitely generated nilpotent group of class 2 with cyclic commutator subgroup.

We introduce the following definition:

Definition. Let G be a finitely generated nilpotent group of class 2. Then $G/Z(G)$ is finitely generated abelian group and thus $G/Z(G) = \langle x_1Z(G) \rangle \times \dots \times \langle x_kZ(G) \rangle$ for some $x_1, \dots, x_k \in G$. The group G is called **d**-group if the following distributive law holds in G ,

$$[x_1^{\alpha_1} \dots x_k^{\alpha_k}, G] = [x_1, G]^{\alpha_1} \dots [x_k, G]^{\alpha_k}$$

where $\alpha_i \in \mathbb{Z}$ and $1 \leq i \leq k$.

Let G be a 2-generator nilpotent group of class 2. It is straightforward to show that G is a **d**-group.

To give an example of an infinite **d**-group, consider the group G with the following presentation

$$G = \langle x_1, x_2, x_3, x_4, x: [x_i, x_j] = x^{m_{ij}}, [x_i, x] = 1; 1 \leq i \leq 4 \text{ and } i < j \rangle,$$

where $m_{i,i+1} = 1$ for all $1 \leq i < 4$ and $m_{ij} = 0$ for all $i + 1 < j$. Then $G' = Z(G) = \langle x \rangle \simeq \mathbb{Z}$ and $G/Z(G) = \langle \bar{x}_1, \bar{x}_2, \bar{x}_3, \bar{x}_4 \rangle \simeq \mathbb{Z}^4$. A quick calculation shows that

$$[x_1^{\alpha_1} x_2^{\alpha_2} x_3^{\alpha_3} x_4^{\alpha_4}, G] = [x_1, G]^{\alpha_1} [x_2, G]^{\alpha_2} [x_3, G]^{\alpha_3} [x_4, G]^{\alpha_4} = \langle x^\alpha \rangle,$$

Where $\alpha_i \in \mathbb{Z}$ for all $1 \leq i \leq 4$ and $\alpha = \text{gcd}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. Therefore G is an infinite **d**-

group.

Now we give a nilpotent group G of class 2 which is not a d -group.

Let G be a free nilpotent group of class 2 on 4 generators a_1, a_2, a_3 and a_4 . If $c_{ij} = [a_i, a_j]$ for $1 \leq i < j \leq 4$, then the relations in G are $[c_{ij}, a_k] = 1$ for $1 \leq i < j \leq 4$ and $1 \leq k \leq 4$, and their consequences. Macdonald in [6] proved that $c_{13}c_{24}$ is not a commutator. Therefore G is not a d -group.

Theorem 1. Let G be a finitely generated nilpotent group of class 2 and

$$G/Z(G) = \langle \bar{x}_1 \rangle \times \dots \times \langle \bar{x}_k \rangle.$$

- (i) There exists a monomorphism $\text{Aut}_{\text{pwi}}(G) \hookrightarrow \prod_{i=1}^k \text{Hom}(\langle \bar{x}_i \rangle, [x_i, G])$.
- (ii) If $[x_i, G]$ is cyclic for all $1 \leq i \leq k$, then there exists a monomorphism $\text{Aut}_{\text{pwi}}(G) \hookrightarrow \text{Inn}(G)$.

In particular if G is a d -group of class 2 then the monomorphisms in (i) and (ii) are isomorphism. Furthermore $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$ if and only if $[x_i, G]$ is cyclic for all $1 \leq i \leq k$.

Notice that if G is a finite group then, as consequence of this result, we derive Theorem 3.5 and Corollary 3.6 of Yadav in [12].

In particular, we derive the following consequences of Theorem 1.

Corollary 1. Let G be a finitely generated nilpotent group of class 2 in which G' is cyclic, then $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$. In particular if G' is finite, then $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$.

Recall that by Corollary 3.6 in [12], in a finite nilpotent group of class 2, if G' is cyclic then $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$. But we cannot hope for a similar conclusion when G is not finite. We will provide an example in the section 2. However, in a finitely generated nilpotent group of class 2, by Corollary 1 we have $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$. So the structure of $\text{Aut}_{\text{pwi}}(G)$ is determined.

Corollary 2. Let G be a finitely generated nilpotent group of class 2. If the commutator subgroup of G is cyclic, then $\text{Aut}_{\text{pwi}}(G)/\text{Inn}(G)$ is torsion.

Let G be a group and N be a non-trivial proper normal subgroup of G . The pair

(G, N) is called a Camina pair if $xN \subseteq x^G$ for all $x \in G \setminus N$. A group G is called a Camina group if (G, G') is a Camina pair.

Clearly, if G is a Camina group of class 2 then it is a d-group. So, as an immediate consequence of Theorem 1, one readily gets the following corollary.

Corollary 3. Let G be a finitely generated nilpotent group of class 2. If G is a Camina group then $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$ if and only if G' is cyclic. Particularly, if $G/Z(G)$ is finite, then $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$ if and only if G' is cyclic.

Preliminary results

Our notation is standard. Let G be a group, by C_m, G' and $Z(G)$, we denote the cyclic group of order m , the commutator subgroup and the center of G , respectively.

If $x, y \in G$, then x^y denotes the conjugate element $y^{-1}xy \in G$. For $x \in G$, x^G denotes the conjugacy class of x in G . The commutator of two elements $x, y \in G$ is defined by $[x, y] = x^{-1}y^{-1}xy$ and more generally, the left-normed commutator of n elements x_1, \dots, x_n is defined inductively by

$$[x_1, \dots, x_{n-1}, x_n] = [x_1, \dots, x_{n-1}]^{-1}x_n^{-1}[x_1, \dots, x_{n-1}]x_n.$$

If $H \leq G$, $[x, H]$ denotes the set of all $[x, h]$ for $h \in H$, this is a subgroup of G when G is of class 2. For any group H and abelian group K , $\text{Hom}(H, K)$ denotes the group of all homomorphisms from H to K . Also C^* is the set of all central automorphisms of G fixing $Z(G)$ elementwise.

Yadav in [12] shows that in a finite nilpotent group of class 2, there exists a monomorphism from $\text{Aut}_{\text{pwi}}(G)$ into $\text{Hom}(G/Z(G), G')$. It turns out that this result remains true when G is an infinite nilpotent group of class 2.

For that, let G be a nilpotent group (finite or infinite) of class 2. Let $\alpha \in \text{Aut}_{\text{pwi}}(G)$. Then the map $g \mapsto g^{-1}\alpha(g)$ is a homomorphism from G into G' . This homomorphism sends $Z(G)$ to 1. So it induces a homomorphism $f_\alpha: G/Z(G) \rightarrow G'$, sending $\bar{g} = gZ(G)$ to $g^{-1}\alpha(g)$, for any $g \in G$. Define

$$\text{Hom}_{\text{pwi}}(G/Z(G), G') = \{f \in \text{Hom}\left(\frac{G}{Z(G)}, G'\right) : f(\bar{g}) \in [g, G] \text{ for all } g \in G\}.$$

To prove $\text{Aut}_{\text{pwi}}(G) \simeq \text{Hom}_{\text{pwi}}(G/Z(G), G')$, we use the following well-known result.

Lemma 1.1 Let N be a normal subgroup of a group G . Let θ be an endomorphism of G such that $\theta(N) \leq N$. Denote by $\bar{\theta}$ and θ_0 the endomorphisms induced by θ in G/N and N , respectively. If $\bar{\theta}$ and θ_0 are surjective (injective), then so is θ .

Proposition 1.2 Let G be a nilpotent group of class 2. Then the above map $\varphi: \alpha \mapsto f_\alpha$ is an isomorphism from $\text{Aut}_{\text{pwi}}(G)$ into $\text{Hom}_{\text{pwi}}(G/Z(G), G')$.

Proof. Since for any $\alpha \in \text{Aut}_{\text{pwi}}(G)$, by the definition $f_\alpha \in \text{Hom}_{\text{pwi}}(G/Z(G), G')$, φ is well defined. Let $\alpha_1, \alpha_2 \in \text{Aut}_{\text{pwi}}(G)$ and $g \in G$. We have $\alpha_1(g^{-1}\alpha_2(g)) = g^{-1}\alpha_2(g)$, since $g^{-1}\alpha_2(g) \in G' \leq Z(G)$. This implies that

$$\begin{aligned} f_{\alpha_1\alpha_2}(\bar{g}) &= g^{-1}\alpha_1(\alpha_2(g)) = g^{-1}\alpha_1(gg^{-1}\alpha_2(g)) \\ &= g^{-1}\alpha_1(g) \cdot g^{-1}\alpha_2(g) = f_{\alpha_1}(\bar{g}) \cdot f_{\alpha_2}(\bar{g}). \end{aligned}$$

Hence φ is a homomorphism. Clearly, φ is injective. Now it suffices to show that φ is surjective.

Let f be any element of $\text{Hom}_{\text{pwi}}(G/Z(G), G')$. By Lemma 1.1 a quick calculation shows that $\varphi(\alpha) = f$, where α is an element of $\text{Aut}_{\text{pwi}}(G)$, sending $g \in G$ to $gf(gZ(G))$. Then we have $\text{Aut}_{\text{pwi}}(G) \simeq \text{Hom}_{\text{pwi}}(G/Z(G), G')$.

* Note that if G is a nilpotent group of class 2 then $\text{Aut}_{\text{pwi}}(G) \simeq \text{Hom}_{\text{pwi}}(G/Z(G), G')$.

It is easy to see that in a Camina nilpotent group of class 2, $\text{Hom}_{\text{pwi}}(G/Z(G), G') = \text{Hom}(G/Z(G), G')$. Hence if G is a Camina group of class 2, then $\text{Aut}_{\text{pwi}}(G) \simeq \text{Hom}(G/Z(G), G')$.

The following well-known facts will be used repeatedly.

Lemma 1.3 Let A, B and C be abelian groups.

- (i) $\text{Hom}(A \times B, C) \simeq \text{Hom}(A, C) \times \text{Hom}(B, C)$.
- (ii) $\text{Hom}(A, B \times C) \simeq \text{Hom}(A, B) \times \text{Hom}(A, C)$.
- (iii) $\text{Hom}(C_m, C_n) \simeq C_d$ where $d = \text{gcd}(m, n)$.
- (iv) $\text{Hom}(\mathbb{Z}, A) \simeq A$.
- (v) If A is torsion group and B is torsion-free group, then $\text{Hom}(A, B) = 1$.
- (vi) If $\text{gcd}(|A|, |B|) \neq 1$, then $\text{Hom}(A, B) \neq 1$.

Main Result

Let G be a finite abelian group. We denote by G_p , the p -primary component of G . Hence $G = \prod_{p \in \pi(G)} G_p$ where $\pi(G)$ denotes the set of all primes p dividing $|G|$.

To prove Theorem 1, we need the following Lemma.

Lemma 2.1 ([1, Corollary 1.4]) Let A and B be two finite abelian groups and $\exp(A) | \exp(B)$. Then $\text{Hom}(A, B) \simeq A$ if and only if $B \simeq C_m \times H$ in which $C_m \simeq \prod_{p \in \pi(A)} B_p$ and $H \simeq \prod_{p \notin \pi(A)} B_p$. In particular, if $\pi(A) = \pi(B)$ then this is equivalent to B is a cyclic group.

Let G be a finitely generated nilpotent group of class 2. Then $G/Z(G)$ is finitely generated abelian group and thus $G/Z(G) = \langle x_1 Z(G) \rangle \times \dots \times \langle x_k Z(G) \rangle$ for some $x_1, \dots, x_k \in G$.

Let $f \in \text{Hom}_{pwi}(G/Z(G), G')$. So $f(gZ(G)) \in [g, G]$ for all $g \in G$. In particular, for all $1 \leq i \leq k$ we have $f(x_i Z(G)) \in [x_i, G]$. Now we prove Theorem 1.

Proof of Theorem 1.

(i) By Proposition 1.2, we have $\text{Aut}_{pwi}(G) \simeq \text{Hom}_{pwi}(G/Z(G), G')$. It suffices to show that there exists a monomorphism from $\text{Hom}_{pwi}(G/Z(G), G')$ into $\prod_{i=1}^k \text{Hom}(\langle \overline{x_i} \rangle, [x_i, G])$. Let $f \in \text{Hom}_{pwi}(G/Z(G), G')$. Denote by f_i , the homomorphism induced by f in $\langle \overline{x_i} \rangle$, for all $1 \leq i \leq k$. Since G is a nilpotent group of class 2, we have $[a^m, b] = [a, b]^m = [a, b^m]$ for each $a, b \in G$ and $m \in \mathbb{Z}$. Consequently, $[x_i^m, G] \leq [x_i, G]$ for all $m \in \mathbb{Z}$ and $1 \leq i \leq k$. Therefore $f_i \in \text{Hom}(\langle \overline{x_i} \rangle, [x_i, G])$. Thus the map α sending any $f \in \text{Hom}_{pwi}(G/Z(G), G')$ to $\alpha(f) = (f_1, \dots, f_k) \in \prod_{i=1}^k \text{Hom}(\langle \overline{x_i} \rangle, [x_i, G])$ is well defined. Now we prove that this map is a monomorphism. Since $(fg)_i = f_i g_i$ for each $f, g \in \text{Hom}_{pwi}(G/Z(G), G')$ and $1 \leq i \leq k$, α is homomorphism. Clearly, $\ker \alpha$ is trivial, this implies that α is monomorphism. Hence the proof of (i) is complete.

(ii) First we show that $[x_i, G]$ is finite if and only if $\langle \overline{x_i} \rangle$ is finite, and further

$\exp([x_i, G]) = \exp(\langle \bar{x}_i \rangle) = |\bar{x}_i|$. For this, let $|[x_i, G]| = n$. Since G is a nilpotent group of class 2, we have $[x_i^n, g] = [x_i, g]^n = 1$ for all $g \in G$ and so $x_i^n \in Z(G)$. Hence $\langle \bar{x}_i \rangle$ is finite and $|\bar{x}_i| |n$. Conversely if $|\bar{x}_i| = m$ then $x_i^m \in Z(G)$ and $[x_i, G]^m = [x_i^m, G] = 1$. Consequently $[x_i, G]$ is finite and $\exp([x_i, G]) = n|m$. Therefore in this case, $m = n$. Hence by Lemma 2.1, for all $1 \leq i \leq k$ we have $\text{Hom}(\langle \bar{x}_i \rangle, [x_i, G]) \simeq \langle \bar{x}_i \rangle$ if and only if $[x_i, G]$ is cyclic.

Now from (i), we have a monomorphism from $\text{Aut}_{\text{pwi}}(G)$ into $\prod_{i=1}^k \text{Hom}(\langle \bar{x}_i \rangle, [x_i, G])$ and therefore we conclude that there exists a monomorphism $\text{Aut}_{\text{pwi}}(G) \hookrightarrow G/Z(G)$, this completes the proof of (ii).

If G is a d -group, then it is easy to see that the monomorphism defined in (i) is an isomorphism from $\text{Aut}_{\text{pwi}}(G)$ into $\prod_{i=1}^k \text{Hom}(\langle \bar{x}_i \rangle, [x_i, G])$.

Finally to complete the proof, it is sufficient to show that if $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$, then $[x_i, G]$ is cyclic for all $1 \leq i \leq k$. Since $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$, by Proposition 1.2 we have $G/Z(G) \simeq \text{Hom}_{\text{pwi}}(G/Z(G), G')$. On the other hand, G is a d -group and hence

$$\text{Hom}_{\text{pwi}}(G/Z(G), G') \simeq \prod_{i=1}^k \text{Hom}(\langle \bar{x}_i \rangle, [x_i, G]).$$

It follows that

$$G/Z(G) = \langle \bar{x}_1 \rangle \times \dots \times \langle \bar{x}_k \rangle \simeq \prod_{i=1}^k \text{Hom}(\langle \bar{x}_i \rangle, [x_i, G]).$$

Now we may assume that $\langle \bar{x}_1 \rangle \times \dots \times \langle \bar{x}_n \rangle$ is the torsion part and $\langle \bar{x}_{n+1} \rangle \times \dots \times \langle \bar{x}_k \rangle$ is the torsion-free part of $G/Z(G)$. Since for all $1 \leq i \leq n$, $\exp([x_i, G]) = \exp(\bar{x}_i) = |\bar{x}_i|$ and $\prod_{i=1}^n \text{Hom}(\langle \bar{x}_i \rangle, [x_i, G]) \simeq \langle \bar{x}_1 \rangle \times \dots \times \langle \bar{x}_n \rangle$, $\text{Hom}(\langle \bar{x}_i \rangle, [x_i, G]) \simeq \langle \bar{x}_i \rangle$ for all $1 \leq i \leq n$ and hence $[x_i, G]$ is cyclic. Furthermore, we have

$$\prod_{i=n+1}^k \text{Hom}(\langle \bar{x}_i \rangle, [x_i, G]) \simeq \langle \bar{x}_{n+1} \rangle \times \dots \times \langle \bar{x}_k \rangle \simeq \mathbb{Z}^{k-n}.$$

Now we have $\text{Hom}(\langle \bar{x}_i \rangle, [x_i, G]) \simeq [x_i, G]$, since $\langle \bar{x}_i \rangle \simeq \mathbb{Z}$ and hence $\prod_{i=n+1}^k [x_i, G] \simeq \mathbb{Z}^{k-n}$. That is $[x_i, G] \simeq \mathbb{Z}$ for all $n + 1 \leq i \leq k$. This implies that $[x_i, G]$ is cyclic for all $1 \leq i \leq k$, as required.

*Notice that if G is a finite group then, as a consequence of this result, we derive Theorem 3.5 and Corollary 3.6 of Yadav in [12].

The following corollary is an easy consequence of the above theorem.

Corollary 2.2 Let G be a finitely generated nilpotent group of class 2 with cyclic commutator subgroup. Then there exists a monomorphism from $\text{Aut}_{\text{pwi}}(G)$ into $\text{Inn}(G)$ or equivalently $\text{Aut}_{\text{pwi}}(G)$ is isomorphic to a subgroup of $G/Z(G)$.

Remark 2.3 We keep here the notation used in Theorem 1.

- (i) By the discussion of (ii) in Theorem 1, if G' is finite cyclic, then $G/Z(G)$ is finite and $|\text{Aut}_{\text{pwi}}(G)| \leq |\text{Inn}(G)| = |G/Z(G)|$. On the other hand, $\text{Inn}(G) \leq \text{Aut}_{\text{pwi}}(G)$ conclude that $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$. Note that in this case, G is not necessarily finite.
- (ii) If G' is infinite cyclic, it follows from the discussion of (ii) in Theorem 1, that $G/Z(G)$ is a free abelian group of finite rank, say $r(G/Z(G)) = k$. We certainly have $\text{Inn}(G) \leq \text{Aut}_{\text{pwi}}(G)$ and thus $r(\text{Inn}(G)) \leq r(\text{Aut}_{\text{pwi}}(G))$. Also $r(\text{Aut}_{\text{pwi}}(G)) \leq r(\text{Inn}(G))$, since $\text{Aut}_{\text{pwi}}(G)$ is isomorphic to a subgroup of $\text{Inn}(G)$. Therefore $\text{Aut}_{\text{pwi}}(G)$ and $\text{Inn}(G)$ have the same rank and hence $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$.

Now it is easy to deduce Corollary 1 from Remark 2.3.

Remark 2.4 It is known that in a nilpotent groups of class 2, $\text{Inn}(G) \leq \text{Aut}_{\text{pwi}}(G) \leq C^*$. So $\text{Inn}(G) = \text{Aut}_{\text{pwi}}(G)$ when $\text{Inn}(G) = C^*$. In [1] we characterized all non torsion-free finitely generated groups in which $\text{Inn}(G) = C^*$. We proved that $\text{Inn}(G) = C^*$ if and only if G is an abelian group or nilpotent of class 2 and $Z(G) \simeq C_m \times H \times \square^r$ in which $C_m \simeq \prod_{p \in \pi(G/Z(G))} Z(G)_p$, $H \simeq \prod_{p \notin \pi(G/Z(G))} Z(G)_p$ and $r \geq 0$ is the torsion-free rank of $Z(G)$ and $G/Z(G)$ has finite exponent.

Hence if G is nilpotent group of class 2, $Z(G) \simeq C_m \times H \times \square^r$ and $G/Z(G)$ has finite exponent then we have $\text{Inn}(G) = \text{Aut}_{\text{pwi}}(G)$. Notice that in this case, G' is cyclic and the equality $\text{Inn}(G) = \text{Aut}_{\text{pwi}}(G)$ also follows from Corollary 1.

Recall that by Corollary 3.6 in [12], in a finite nilpotent group of class 2, if G' is cyclic then $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$. But we cannot hope for a similar conclusion when G is not finite.

For example, consider countably infinitely many copies H_1, H_2, \dots of a given nilpotent group H of class 2 with cyclic commutator subgroup. Let G (respectively, \bar{G}) be the direct product (the cartesian product) of the family $(H_i)_{i>0}$. Clearly, G and \bar{G} are nilpotent of class 2. For each integer $i > 0$, choose an element $a_i \in H_i$ which is not in the center of H_i . Then the inner automorphism of \bar{G} defined by $\bar{\alpha}((t_i)_{i>0}) = (a_i^{-1} t_i a_i)_{i>0}$ induces in G a pointwise inner automorphism α which is not inner (see [4]).

However, in a finitely generated nilpotent group of class 2 with cyclic commutator subgroup, we have $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$, by Corollary 1. So the structure of $\text{Aut}_{\text{pwi}}(G)$ is determined.

Furthermore it is fairly easy to deduce Corollary 2 from Remark 2.3.

We end this part of the paper with some examples of infinite groups G satisfying the conditions of Corollary 1 and therefore $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$.

Example 2.5 Let $G = \langle x_1, x_2, y_1, y_2 : x_1^p = x_2^p = y_1^p = 1, [x_1, x_2] = y_1, [y_1, y_2] = [x_i, y_j] = 1; 1 \leq i, j \leq 2 \rangle$. Then G satisfies the condition of Corollary 1. We have $G' = \langle y_1 \rangle \simeq C_p$, $Z(G) = \langle y_1, y_2 \rangle \simeq C_p \times \mathbb{Z}$ and $G/Z(G) = \langle \bar{x}_1, \bar{x}_2 \rangle \simeq C_p \times C_p$ and hence $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$.

Example 2.6 Let $G = \langle x_1, x_2, x : [x_1, x_2] = x, [x_i, x] = 1; 1 \leq i \leq 2 \rangle$. Then G satisfies the condition of Corollary 1. We have $G' = Z(G) = \langle x \rangle \simeq \mathbb{Z}$ and $\frac{G}{Z(G)} = \langle \bar{x}_1, \bar{x}_2 \rangle \simeq \mathbb{Z} \times \mathbb{Z}$. Hence $\text{Aut}_{\text{pwi}}(G) \simeq \text{Inn}(G)$. It is easy to see that in this case every pointwise inner automorphism is inner and so $\text{Aut}_{\text{pwi}}(G) = \text{Inn}(G)$ (see [1, Example 3.4]).

Acknowledgements

We thank the editor of the Journal Science of Kharazmi University and the referees who patiently read and verified this paper.

References

1. Z. Azhdari, M. Akhavan-Malayeri, "On Inner automorphisms and central automorphisms of

- Nilpotent Group of Class 2", J. Algebra Appl. Vol. 10, No. 4 (2011) 1283-1290.
2. W. Burnside, "Theory of groups of finite order, 2nd Ed. Dover Publications", Inc., 1955. Reprint of the 2nd edition (Cambridge, 1911).
 3. W. Burnside, "On the outer automorphisms of a group", Proc. London Math. Soc. (2) 11 (1913) 40-42.
 4. G. Endimioni, "Pointwise inner automorphism in a free nilpotent group", Quart. J. Math. 53 (2002) 397-402.
 5. E. K. Grossman, "On the residual finiteness of certain mapping class group", J. London Math. Soc. (Ser. 2) 9 (1974/75) 160-164.
 6. I. D. Macdonald, "On a set of normal subgroups", Proc. Glasgow Math. Soc. 5 (1962) 137-146.
 7. T. Ono and H. Wada, "Hasse principle" for symmetric and alternating groups, Proc. Japan Acad., 75 Ser. A (1999) 61-62.
 8. C. H. Sah, "Automorphisms of finite groups", Journal of Algebra 10 (1968) 47-68.
 9. D. Segal, "On the outer automorphism group of a polycyclic group", in Proceedings Bressanone/Brixen 1989, Suppl. ai Rend. Del Circ. Math. di Palermo 23, Palermo (1990) 265-278.
 10. H. Wada, "Hasse principle" for $SL_n(D)$, Proc. Japan Acad. 75 Ser. A (1999) 67-69.
 11. H. Wada, "Hasse principle" for $GL_n(D)$, Proc. Japan Acad. 76 Ser. A (2000) 44-46.
 12. M. K. Yadav, "On automorphisms of some finite p-groups", Proc. Indian Acad. Sci. (Math. Sci) 118(1) (2008) 1-11.